



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2011, 22nd Annual JWP Conference

Apr 9th, 2:35 PM - 3:35 PM

Evaluation of DOM Uptake by Spionid Polychaete Larvae from Antarctica

Travis Rosenkranz

Illinois Wesleyan University

William Jaeckle, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

 Part of the [Biology Commons](#)

Travis Rosenkranz and William Jaeckle, Faculty Advisor, "Evaluation of DOM Uptake by Spionid Polychaete Larvae from Antarctica" (April 9, 2011). *John Wesley Powell Student Research Conference*. Paper 19.
<http://digitalcommons.iwu.edu/jwprc/2011/posters2/19>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P40

**EVALUATION OF DOM UPTAKE BY SPIONID POLYCHAETE LARVAE
FROM ANTARCTICA**

Travis Rosenkranz and William Jaeckle*
Biology Department, Illinois Wesleyan University

Thorson (1946, 1950) hypothesized that polar invertebrate species avoid a pelagic larval phase and instead favor direct development (“Thorson’s Rule”) and proposed an inverse relationship existed between latitude and the number of species with planktonic larval stages. Recent studies, however, suggest that a decrease in invertebrate larval abundance, not diversity, is correlated with an increase in latitude. The oceans surrounding Antarctica are nutrient rich, especially during late spring and early summer. Larvae of Antarctic spionid polychaetes feed using a dual-band system of cilia whose activities concentrate and then capture food particles. We evaluated the ability of spionid larvae from Antarctica to process seawater and absorb dissolved organic materials (DOM). Collected larvae were exposed to the iron-containing polysaccharide iron-dextran (1 mg / mL) for 12 to 24 hours at 0 °C. The presence of the iron (from iron-dextran) in larval tissue was detected using the “Prussian Blue” reaction. We found that the label was localized solely in the digestive system; there was no evidence of uptake by the epidermis. Control larvae, not exposed to iron-dextran, showed no label in tissue. Antarctic spionid larvae transport seawater through their digestive system and can potentially exploit DOM as an alternate source of food.